

To appear in:
Rend. Sem. Mat. Univ. Pol. Torino (1997)

THE JACOBI FLOW

PETER W. MICHOR

Erwin Schrödinger Institut für Mathematische
Physik, Pasteurgasse 6/7, A-1090 Wien, Austria

February 1, 2008

For Wlodek Tulczyjew, on the occasion of his 65th birthday.

It is well known that the geodesic flow on the tangent bundle is the flow of a certain vector field which is called the spray $S : TM \rightarrow TTM$. It is maybe less well known that the flow lines of the vector field $\kappa_{TM} \circ TS : TTM \rightarrow TTTM$ project to Jacobi fields on TM . This could be called the ‘Jacobi flow’. This result was developed for the lecture course [5], and it is the main result of this paper. I was motivated by the paper [6] of Urbanski in these proceedings to publish it, as an explanation of some of the uses of iterated tangent bundles in differential geometry.

1. The tangent bundle of a vector bundle. Let (E, p, M) be a vector bundle with fiber addition $+_E : E \times_M E \rightarrow E$ and fiber scalar multiplication $m_t^E : E \rightarrow E$. Then (TE, π_E, E) , the tangent bundle of the manifold E , is itself a vector bundle, with fiber addition denoted by $+_{TE}$ and scalar multiplication denoted by m_t^{TE} .

If $(U_\alpha, \psi_\alpha : E \upharpoonright U_\alpha \rightarrow U_\alpha \times V)_{\alpha \in A}$ is a vector bundle atlas for E , such that (U_α, u_α) is a manifold atlas for M , then $(E \upharpoonright U_\alpha, \psi'_\alpha)_{\alpha \in A}$ is an atlas for the manifold E , where

$$\psi'_\alpha := (u_\alpha \times \text{Id}_V) \circ \psi_\alpha : E \upharpoonright U_\alpha \rightarrow U_\alpha \times V \rightarrow u_\alpha(U_\alpha) \times V \subset \mathbb{R}^m \times V.$$

Hence the family $(T(E \upharpoonright U_\alpha), T\psi'_\alpha : T(E \upharpoonright U_\alpha) \rightarrow T(u_\alpha(U_\alpha) \times V) = u_\alpha(U_\alpha) \times V \times \mathbb{R}^m \times V)_{\alpha \in A}$ is the atlas describing the canonical vector bundle structure of (TE, π_E, E) . The transition functions are in turn:

$$\begin{aligned} (\psi_\alpha \circ \psi_\beta^{-1})(x, v) &= (x, \psi_{\alpha\beta}(x)v) \quad \text{for } x \in U_{\alpha\beta} \\ (u_\alpha \circ u_\beta^{-1})(y) &= u_{\alpha\beta}(y) \quad \text{for } y \in u_\beta(U_{\alpha\beta}) \\ (\psi'_\alpha \circ (\psi'_\beta)^{-1})(y, v) &= (u_{\alpha\beta}(y), \psi_{\alpha\beta}(u_\beta^{-1}(y))v) \\ (T\psi'_\alpha \circ T(\psi'_\beta)^{-1})(y, v; \xi, w) &= (u_{\alpha\beta}(y), \psi_{\alpha\beta}(u_\beta^{-1}(y))v; d(u_{\alpha\beta})(y)\xi, \\ &\quad (d(\psi_{\alpha\beta} \circ u_\beta^{-1})(y))\xi)v + \psi_{\alpha\beta}(u_\beta^{-1}(y))w). \end{aligned}$$

1991 *Mathematics Subject Classification.* 53C22.

Key words and phrases. Spray, geodesic flow, Jacobi flow, higher tangent bundles.

Supported by ‘Fonds zur Förderung der wissenschaftlichen Forschung, Projekt P 10037 PHY’.

So we see that for fixed (y, v) the transition functions are linear in $(\xi, w) \in \mathbb{R}^m \times V$. This describes the vector bundle structure of the tangent bundle (TE, π_E, E) .

For fixed (y, ξ) the transition functions of TE are also linear in $(v, w) \in V \times V$. This gives a vector bundle structure on (TE, Tp, TM) . Its fiber addition will be denoted by $T(+_E) : T(E \times_M E) = TE \times_{TM} TE \rightarrow TE$, since it is the tangent mapping of $+_E$. Likewise its scalar multiplication will be denoted by $T(m_t^E)$. One may say that the second vector bundle structure on TE , that one over TM , is the derivative of the original one on E .

The space $\{\Xi \in TE : Tp.\Xi = 0 \text{ in } TM\} = (Tp)^{-1}(0)$ is denoted by VE and is called the *vertical bundle* over E . The local form of a vertical vector Ξ is $T\psi'_\alpha.\Xi = (y, v; 0, w)$, so the transition functions are $(T\psi'_\alpha \circ T(\psi'_\beta)^{-1})(y, v; 0, w) = (u_{\alpha\beta}(y), \psi_{\alpha\beta}(u_\beta^{-1}(y))v; 0, \psi_{\alpha\beta}(u_\beta^{-1}(y))w)$. They are linear in $(v, w) \in V \times V$ for fixed y , so VE is a vector bundle over M . It coincides with $0_M^*(TE, Tp, TM)$, the pullback of the bundle $TE \rightarrow TM$ over the zero section. We have a canonical isomorphism $Vl_E : E \times_M E \rightarrow VE$, called the *big vertical lift*, given by $Vl_E(u_x, v_x) := \partial_t|_0(u_x + tv_x)$, which is fiber linear over M . We will mainly use the *small vertical lift* $vl_E : E \rightarrow TE$, given by $vl_E(v_x) = \partial_t|_0 t.v_x = Vl_E(0_x, v_x)$. The local representation of the vertical lift is $(T\psi'_\alpha \circ vl_E \circ (\psi'_\alpha)^{-1})(y, v) = (y, 0; 0, v)$.

If $\varphi : (E, p, M) \rightarrow (F, q, N)$ is a vector bundle homomorphism, then we have $vl_F \circ \varphi = T\varphi \circ vl_E : E \rightarrow VF \subset TF$. So vl is a natural transformation between certain functors on the category of vector bundles and their homomorphisms. The mapping $vrp_E := pr_2 \circ Vl_E^{-1} : VE \rightarrow E$ is called the *vertical projection*.

2. The second tangent bundle of a manifold. All of 1 is valid for the second tangent bundle TTM of a manifold, but here we have one more natural structure at our disposal. The *canonical flip* or *involution* $\kappa_M : TTM \rightarrow TTM$ is defined locally by

$$(TTu \circ \kappa_M \circ TTu^{-1})(x, \xi; \eta, \zeta) = (x, \eta; \xi, \zeta),$$

where (U, u) is a chart on M . Clearly this definition is invariant under changes of charts (Tu_α equals ψ'_α from 1).

The flip κ_M has the following properties:

- (1) $\kappa_N \circ TTf = TTf \circ \kappa_M$ for each $f \in C^\infty(M, N)$.
- (2) $T(\pi_M) \circ \kappa_M = \pi_{TM}$ and $\pi_{TM} \circ \kappa_M = T(\pi_M)$.
- (3) $\kappa_M^{-1} = \kappa_M$.
- (4) κ_M is a linear isomorphism from the vector bundle $(TTM, T(\pi_M), TM)$ to the bundle (TTM, π_{TM}, TM) , so it interchanges the two vector bundle structures on TTM .
- (5) It is the unique smooth mapping $TTM \rightarrow TTM$ which satisfies

$$\partial_t \partial_s c(t, s) = \kappa_M \partial_s \partial_t c(t, s)$$

for each $c : \mathbb{R}^2 \rightarrow M$.

All this follows from the local formula given above. A quite early use of κ_M is in [4].

3. Lemma. For vector fields $X, Y \in \mathfrak{X}(M)$ we have

$$\begin{aligned} [X, Y] &= \text{vrp}_{TM} \circ (TY \circ X - \kappa_M \circ TX \circ Y), \\ TY \circ X - {}_{TM} \kappa_T \circ TX \circ Y &= \text{vl}_{TM}(Y, [X, Y]) \\ &= (\text{vl}_{TM} \circ [X, Y]) T(+_{TM}) (0_{TM} \circ Y). \end{aligned}$$

See [3] 6.13, 6.19, or 37.13 for different proofs of this well known result.

4. Linear connections and their curvatures. Let (E, p, M) be a vector bundle. Recall that a linear connection on the vector bundle E can be described by specifying its *connector* $K : TE \rightarrow E$. This notions seems to be due to [2]. Any smooth mapping $K : TE \rightarrow E$ which is a (fiber linear) homomorphism for both vector bundle structures on TE ,

$$\begin{array}{ccc} TE & \xrightarrow{K} & E \\ \pi_E \downarrow & & \downarrow p \\ E & \xrightarrow{p} & M \end{array} \quad \begin{array}{ccc} TE & \xrightarrow{K} & E \\ Tp \downarrow & & \downarrow p \\ TM & \xrightarrow{\pi_M} & M \end{array}$$

and which is a left inverse to the vertical lift, $K \circ \text{vl}_E = \text{Id}_E : E \rightarrow TE \rightarrow E$, specifies a linear connection. Namely: The inverse image $H := K^{-1}(0_E)$ of the zero section $0_E \subset E$, it is a subvector bundle for both vector bundle structures, and for the vector bundle structure $\pi_E : TE \rightarrow E$ the subbundle H turns out to be a complementary bundle for the vertical bundle $VE \rightarrow E$. We get then the associated *horizontal lift mapping*

$$C : TM \times_M E \rightarrow TE, \quad C(\cdot, u) = \left(Tp|_{\ker(K : T_u E \rightarrow E_{p(u)})} \right)^{-1}$$

which has the following properties

$$\begin{aligned} (Tp, \pi_E) \circ C &= \text{Id}_{TM \times_M E}, \\ C(\cdot, u) : T_{p(u)} M &\rightarrow T_u E \text{ is linear for each } u \in E, \\ C(X_x, \cdot) : E_x &\rightarrow (Tp)^{-1}(X_x) \text{ is linear for each } X_x \in T_x M. \end{aligned}$$

Conversely given a smooth horizontal lift mapping C with these properties one can reconstruct a connector K .

For any manifold N , smooth mapping $s : N \rightarrow E$ along $f = p \circ s : N \rightarrow M$, and vector field $X \in \mathfrak{X}(N)$ a connector $K : TE \rightarrow E$ defines the *covariant derivative* of s along X by

$$(1) \quad \nabla_X s := K \circ Ts \circ X : N \rightarrow TN \rightarrow TE \rightarrow E.$$

See the following diagram for all the mappings.

$$(2) \quad \begin{array}{ccccc} & & TE & & \\ & Ts \nearrow & \downarrow \pi_E & \searrow K & \\ TN & & E & & E \\ \uparrow X & \nearrow s & \nearrow \nabla_X s & & \\ N & \xrightarrow{f} & M & & \end{array}$$

In canonical coordinates as in 1 we have then

$$\begin{aligned} C((y, \xi), (y, v)) &= (y, v; \xi, \Gamma_y(v, \xi)), \\ K(y, v; \xi, w) &= (y, w - \Gamma_y(v, \xi)), \\ \nabla_{(y, \xi)}(\text{Id}, s) &= (\text{Id}, ds(y)\xi - \Gamma_y(s(y), \xi)), \end{aligned}$$

where the *Christoffel symbol* $\Gamma_y(v, \xi)$ is smooth in y and bilinear in (v, ξ) . Here the sign is the negative of the one in many more traditional approaches, since Γ parametrizes the horizontal bundle.

Let $C_f^\infty(N, E)$ denote the space of all sections along f of E , isomorphic to the space $C^\infty(f^*E)$ of sections of the pullback bundle. The covariant derivative may then be viewed as a bilinear mapping $\nabla : \mathfrak{X}(N) \times C_f^\infty(N, E) \rightarrow C_f^\infty(N, E)$. It has the following properties which follow directly from the definitions:

- (3) $\nabla_X s$ is $C^\infty(N, \mathbb{R})$ -linear in $X \in \mathfrak{X}(N)$. For $x \in N$ also we have $\nabla_{X(x)} s = K.Ts.X(x) = (\nabla_X s)(x) \in E$.
- (4) $\nabla_X(h.s) = dh(X).s + h.\nabla_X s$ for $h \in C^\infty(N, \mathbb{R})$.
- (5) For any manifold Q , smooth mapping $g : Q \rightarrow N$, and $Y_y \in T_y Q$ we have $\nabla_{Tg.Y_y} s = \nabla_{Y_y}(s \circ g)$. If $Y \in \mathfrak{X}(Q)$ and $X \in \mathfrak{X}(N)$ are g -related, then we have $\nabla_Y(s \circ g) = (\nabla_X s) \circ g$.

For vector fields $X, Y \in \mathfrak{X}(M)$ and a section $s \in C^\infty(E)$ the curvature $R \in \Omega^2(M, L(E, E))$ of the connection is given by

$$(6) \quad R(X, Y)s = ([\nabla_X, \nabla_Y] - \nabla_{[X, Y]})s$$

Theorem. Let $K : TE \rightarrow E$ be the connector of a linear connection on a vector bundle (E, p, M) . If $s : N \rightarrow E$ is a section along $f := p \circ s : N \rightarrow M$ then we have for vector fields $X, Y \in \mathfrak{X}(N)$

$$\begin{aligned} (7) \quad \nabla_X \nabla_Y s - \nabla_Y \nabla_X s - \nabla_{[X, Y]} s &= \\ &= (K \circ TK \circ \kappa_E - K \circ TK) \circ TTs \circ TX \circ Y = \\ &= R \circ (Tf \circ X, Tf \circ Y)s : N \rightarrow E, \end{aligned}$$

where $R \in \Omega^2(M; L(E, E))$ is the curvature.

Proof. Let first $m_t^E : E \rightarrow E$ denote the scalar multiplication. Then we have $\partial_t|_0 m_t^E = \text{vl}_E$ where $\text{vl}_E : E \rightarrow TE$ is the vertical lift. We use then lemma 3 and some obvious commutation relations to get in turn:

$$\begin{aligned} \text{vl}_E \circ K &= \partial_t|_0 m_t^E \circ K = \partial_t|_0 K \circ m_t^{TE} = TK \circ \partial_t|_0 m_t^{TE} = TK \circ \text{vl}_{(TE, \pi_E, E)} \cdot \\ \nabla_X \nabla_Y s - \nabla_Y \nabla_X s - \nabla_{[X, Y]} s &= \\ &= K \circ T(K \circ Ts \circ Y) \circ X - K \circ T(K \circ Ts \circ X) \circ Y - K \circ Ts \circ [X, Y] \\ K \circ Ts \circ [X, Y] &= K \circ \text{vl}_E \circ K \circ Ts \circ [X, Y] \\ &= K \circ TK \circ \text{vl}_{TE} \circ Ts \circ [X, Y] = K \circ TK \circ TTs \circ \text{vl}_{TN} \circ [X, Y] \\ &= K \circ TK \circ TTs \circ ((TY \circ X - \kappa_N \circ TX) \circ Y) (T-) 0_{TN} \circ Y \\ &= K \circ TK \circ TTs \circ TY \circ X - K \circ TK \circ TTs \circ \kappa_N \circ TX \circ Y - 0. \end{aligned}$$

Now we sum up and use $TTs \circ \kappa_N = \kappa_E \circ TTs$ to get the first result. If in particular we choose $f = \text{Id}_M$ so that s is a section of $E \rightarrow M$ and X, Y are vector fields on M , then we get the curvature R .

To see that in the general case $(K \circ TK \circ \kappa_E - K \circ TK) \circ TTs \circ TX \circ Y$ coincides with $R(Tf \circ X, Tf \circ Y)s$ one has to write out (1) and $(TTs \circ TX \circ Y)(x) \in TTE$ in canonical charts induced from vector bundle charts of E . $\square$

5. Torsion. Let $K : TTM \rightarrow M$ be a linear connector on the tangent bundle, let $X, Y \in \mathfrak{X}(M)$. Then the torsion is given by

$$\text{Tor}(X, Y) = (K \circ \kappa_M - K) \circ TX \circ Y.$$

If moreover $f : N \rightarrow M$ is smooth and $U, V \in \mathfrak{X}(N)$ then we get also

$$\begin{aligned} \text{Tor}(Tf.U, Tf.V) &= \nabla_U(Tf \circ V) - \nabla_V(Tf \circ U) - Tf \circ [U, V] \\ &= (K \circ \kappa_M - K) \circ TTf \circ TU \circ V. \end{aligned}$$

Proof. (9) We have in turn

$$\begin{aligned} \text{Tor}(X, Y) &= \nabla_X Y - \nabla_Y X - [X, Y] \\ &= K \circ TY \circ X - K \circ TX \circ Y - K \circ \text{vl}_{TM} \circ [X, Y] \\ K \circ \text{vl}_{TM} \circ [X, Y] &= K \circ ((TY \circ X - \kappa_M \circ TX \circ Y) (T-) 0_{TM} \circ Y) \\ &= K \circ TY \circ X - K \circ \kappa_M \circ TX \circ Y - 0. \end{aligned}$$

An analogous computation works in the second case, and that $(K \circ \kappa_M - K) \circ TTf \circ TU \circ V = \text{Tor}(Tf.U, Tf.V)$ can again be checked in local coordinates. $\square$

6. Sprays. Given a linear connector $K : TTM \rightarrow M$ on the tangent bundle with its horizontal lift mapping $C : TM \times_M TM \rightarrow TTM$, then $S := C \circ \text{diag} : TM \rightarrow TM \times_M TM \rightarrow TTM$ is called the *spray*. This notion is due to [1]. The spray has the following properties:

$$\begin{aligned} \pi_{TM} \circ S &= \text{Id}_{TM} && \text{a vector field on } TM, \\ T(\pi_M) \circ S &= \text{Id}_{TM} && \text{a second order differential equation,} \\ S \circ m_t^{TM} &= T(m_t^{TM}) \circ m_t^{TTM} \circ S && \text{'quadratic',} \end{aligned}$$

where m_t^E is the scalar multiplication by t on a vector bundle E . From S one can reconstruct the torsion free part of C . The following result is well known:

Lemma. For a spray $S : TM \rightarrow TTM$ on M , for $X \in TM$

$$\text{geo}^S(X)(t) := \pi_M(\text{Fl}_t^S(X))$$

defines a geodesic structure on M , where Fl^S is the flow of the vector field S .

The abstract properties of a geodesic structure are obvious:

$$\begin{aligned} \text{geo} : TM \times \mathbb{R} \supset U &\rightarrow M \\ \text{geo}(X)(0) &= \pi_M(X), \quad \partial_t|_0 \text{geo}(X)(t) = X \\ \text{geo}(tX)(s) &= \text{geo}(X)(ts) \\ \text{geo}(\text{geo}(X)'(t))(s) &= \text{geo}(X)(t+s) \end{aligned}$$

From a geodesic structure one can reconstruct the spray by differentiation.

7. Theorem. *Let $S : TM \rightarrow TTM$ be a spray on a manifold M . Then $\kappa_{TM} \circ TS : TTM \rightarrow TTTM$ is a vector field. Consider a flow line*

$$Y(t) = \text{Fl}_t^{\kappa_{TM} \circ TS}(Y(0))$$

of this field. Then we have:

- $c := \pi_M \circ \pi_{TM} \circ Y$ is a geodesic on M .
- $\dot{c} = \pi_{TM} \circ Y$ is the velocity field of c .
- $J := T(\pi_M) \circ Y$ is a Jacobi field along c .
- $\dot{J} = \kappa_M \circ Y$ is the velocity field of J .
- $\nabla_{\partial_t} J = K \circ \kappa_M \circ Y$ is the covariant derivative of J .
- The Jacobi equation is given by:

$$\begin{aligned} 0 &= \nabla_{\partial_t} \nabla_{\partial_t} J + R(J, \dot{c})\dot{c} + \nabla_{\partial_t} \text{Tor}(J, \dot{c}) \\ &= K \circ TK \circ TS \circ Y. \end{aligned}$$

This implies that in a canonical chart induced from a chart on M the curve $Y(t)$ is given by

$$(c(t), c'(t); J(t), J'(t)).$$

Proof. Consider a curve $s \mapsto X(s)$ in TM . Then each $t \mapsto \pi_M(\text{Fl}_t^S(X(s)))$ is a geodesic in M , and in the variable s it is a variation through geodesics. Thus $J(t) := \partial_s|_0 \pi_M(\text{Fl}_t^S(X(s)))$ is a Jacobi field along the geodesic $c(t) := \pi_M(\text{Fl}_t^S(X(0)))$, and each Jacobi field is of this form, for a suitable curve $X(s)$. We consider now the curve $Y(t) := \partial_s|_0 \text{Fl}_t^S(X(s))$ in TTM . Then by 2.(6) we have

$$\begin{aligned} \partial_t Y(t) &= \partial_t \partial_s|_0 \text{Fl}_t^S(X(s)) = \kappa_{TM} \partial_s|_0 \partial_t \text{Fl}_t^S(X(s)) = \kappa_{TM} \partial_s|_0 S(\text{Fl}_t^S(X(s))) \\ &= (\kappa_{TM} \circ TS)(\partial_s|_0 \text{Fl}_t^S(X(s))) = (\kappa_{TM} \circ TS)(Y(t)), \end{aligned}$$

so that $Y(t)$ is a flow line of the vector field $\kappa_{TM} \circ TS : TTM \rightarrow TTTM$. Moreover using the properties of κ from section 2 and of S from section 6 we get

$$\begin{aligned} T\pi_M.Y(t) &= T\pi_M.\partial_s|_0 \text{Fl}_t^S(X(s)) = \partial_s|_0 \pi_M(\text{Fl}_t^S(X(s))) = J(t), \\ \pi_M T\pi_M Y(t) &= c(t), \text{ the geodesic,} \\ \partial_t J(t) &= \partial_t T\pi_M.\partial_s|_0 \text{Fl}_t^S(X(s)) = \partial_t \partial_s|_0 \pi_M(\text{Fl}_t^S(X(s))), \\ &= \kappa_M \partial_s|_0 \partial_t \pi_M(\text{Fl}_t^S(X(s))) = \kappa_M \partial_s|_0 \partial_t \pi_M(\text{Fl}_t^S(X(s))) \\ &= \kappa_M \partial_s|_0 T\pi_M.\partial_t \text{Fl}_t^S(X(s)) = \kappa_M \partial_s|_0 (T\pi_M \circ S) \text{Fl}_t^S(X(s)) \\ &= \kappa_M \partial_s|_0 \text{Fl}_t^S(X(s)) = \kappa_M Y(t), \\ \nabla_{\partial_t} J &= K \circ \partial_t J = K \circ \kappa_M \circ Y. \end{aligned}$$

Finally let us express the well known Jacobi expression, where we put $\gamma(t, s) := \pi_M(\text{Fl}_t^S(X(s)))$ for short and use most of the expressions from above:

$$\begin{aligned} \nabla_{\partial_t} \nabla_{\partial_t} J + R(J, \dot{c})\dot{c} + \nabla_{\partial_t} \text{Tor}(J, \dot{c}) &= \\ &= \nabla_{\partial_t} \nabla_{\partial_t} T\gamma.\partial_s + R(T\gamma.\partial_s, T\gamma.\partial_t)T\gamma.\partial_t + \nabla_{\partial_t} \text{Tor}(T\gamma.\partial_s, T\gamma.\partial_t) \\ &= K.T(K.T(T\gamma.\partial_s).\partial_t).\partial_t \\ &\quad + (K.TK.\kappa_{TM} - K.TK).TT(T\gamma.\partial_t).T\partial_s.\partial_t \\ &\quad + K.T((K.\kappa_M - K).TT\gamma.T\partial_s.\partial_t).\partial_t \end{aligned}$$

Note that for example for the term in the second summand we have

$$TTT\gamma.TT\partial_t.T\partial_s.\partial_t = T(T(\partial_t\gamma).\partial_s).\partial_t = \partial_t\partial_s\partial_t\gamma = \partial_t.\kappa_M.\partial_t.\partial_s\gamma = T\kappa_M.\partial_t.\partial_t.\partial_s\gamma$$

which at $s = 0$ equals $T\kappa_M.\ddot{J}$. Using this we get for the Jacobi expression at $s = 0$:

$$\begin{aligned} \nabla_{\partial_t}\nabla_{\partial_t}J + R(J,\dot{c})\dot{c} + \nabla_{\partial_t}\text{Tor}(J,\dot{c}) &= \\ &= (K.TK + K.TK.\kappa_{TM}.T\kappa_M - K.TK.T\kappa_M + K.TK.T\kappa_M - K.TK).\partial_t\partial_tJ = \\ &= K.TK.\kappa_{TM}.T\kappa_M.\partial_t\partial_tJ = K.TK.\kappa_{TM}.\partial_tY = K.TK.TS.Y, \end{aligned}$$

where we used $\partial_t\partial_tJ = \partial_t(\kappa_M.Y) = T\kappa_M\partial_tY = T\kappa_M.\kappa_{TM}.TS.Y$. Finally the validity of the Jacobi equation $0 = K.TK.TS.Y$ follows trivially from $K \circ S = 0_{TM}$. $\square$

REFERENCES

1. Ambrose, W; Palais, R.S., Singer I.M., *Sprays*, Acad. Brasileira de Ciencias **32** (1960), 163–178.
2. Gromoll, D.; Klingenberg, W.; Mayer, W., *Riemannsche Geometrie im Großen*, Lecture Notes in Math. 55, Springer-Verlag, Berlin, Heidelberg, 1968.
3. Kolář, Ivan; Michor, Peter W.; Slovák, Jan, *Natural operations in differential geometry*, Springer-Verlag, Berlin, Heidelberg, New York, 1993.
4. Losik, M.V., *On infinitesimal connections in tangential stratifiable spaces*, Izv. Vyssh. Uchebn. Zaved., Mat. **5(42)** (1964), 54–60. (Russian)
5. Michor, Peter W., *Riemannsche Differentialgeometrie*, Lecture course at the Universität Wien, 1988/89.
6. Urbanski, P., *Double bundles*, These proceedings ?? (??), ??.

INSTITUT FÜR MATHEMATIK, UNIVERSITÄT WIEN, STRUDLHOFGASSE 4, A-1090 WIEN, AUSTRIA

E-mail address: Peter.Michor@esi.ac.at